

ON RAMANUJAN'S CUBIC CONTINUED FRACTION AND EXPLICIT EVALUATIONS OF THETA-FUNCTIONS

CHANDRASHEKRA ADIGA, TAEKYUN KIM,*
M. S. MAHADEVA NAIKA, AND H. S. MADHUSUDHAN

*Department of Studies in Mathematics, University of Mysore
Manasagangotri, MYSORE-570 006, INDIA*

**Institute of Science Education, Kongju National University
Kongju 314-701, REPUBLIC OF KOREA*

ABSTRACT. In this paper we give two integral representations for the Ramanujan's cubic continued fraction $V(q)$ and also derive a modular equation relating $V(q)$ and $V(q^3)$. We also establish some modular equations and a transformation formula for Ramanujan's theta-function. As an application of these, we compute several new explicit evaluations of theta-functions and Ramanujan's cubic continued fraction.

1. Introduction

The celebrated Rogers-Ramanujan continued fraction is defined by

$$R(q) := \frac{q^{1/5}}{1} \frac{q}{+1} \frac{q^2}{+1} \frac{q^3}{+1} \frac{q^4}{+\dots}, \quad |q| < 1. \quad (1.1)$$

On page 46 in his 'lost' notebook [19, p. 46], Ramanujan claims that

$$R(q) = \frac{\sqrt{5}-1}{2} \exp \left((-1/5) \int_q^1 \frac{(1-t)^5(1-t^2)^5 \dots dt}{(1-t^5)(1-t^{10}) \dots t} \right) \quad (1.2)$$

$$= \frac{\sqrt{5}-1}{2} - \frac{\sqrt{5}}{1 + \frac{3+\sqrt{5}}{2}} \exp \left((1/5) \int_0^q \frac{(1-t)^5(1-t^2)^5 \dots dt}{(1-t^{1/5})(1-t^{2/5}) \dots t^{4/5}} \right), \quad (1.3)$$

Key words. Cubic continued fraction, integral representation, modular equation, theta-function, transformation formula.

2000 AMS Subject Classification: 11A55, 33D15, 33D20.

where $0 < q < 1$. The first equality (1.2) was proved by G.E. Andrews [3] and the second equality (1.3) was proved by Seung Hwan Son [20]. On page 207 of his 'lost' notebook Ramanujan also recorded six identities involving integrals of theta-functions. All these identities were proved by S.H. Son [20] and were generalized by Chandrashekar Adiga, K. R. Vasuki and M. S. Mahadeva Naika [1]. On page 365 of his 'lost' notebook [19], Ramanujan wrote five modular equations relating $R(q)$ with $R(-q)$, $R(q^2)$, $R(q^3)$, $R(q^4)$, and $R(q^5)$.

Ramanujan eventually found several generalizations and ramifications of (1.1) which are recorded in his 'lost' notebook. These and related works may be found in the papers by S. Bhargava [8], S. Bhargava and C. Adiga [9], [10], R. Y. Denis [13], [14], [15].

On page 366 of his 'lost' notebook [19], Ramanujan investigated the continued fraction

$$V(q) := \frac{q^{1/3}}{1} \frac{q + q^2}{+ 1} \frac{q^2 + q^4}{+ 1} \frac{q^4 + q^8}{+ \dots}, \quad |q| < 1, \quad (1.4)$$

which is known as Ramanujan's cubic continued fraction. Analogous to those relations for $R(q)$ which are mentioned above, H. H. Chan [12] established several modular equations relating $V(q)$ with $V(-q)$, $V(q^2)$ and $V(q^3)$. An example of these modular equations is

$$V^3(q) = V(q^3) \frac{1 - V(q^3) + V^2(q^3)}{1 + 2V(q^3) + 4V^2(q^3)}. \quad (1.5)$$

In Section 2, we will establish two integral representations for $V(q)$, which are analogous to (1.2) and (1.3). We also give a simple proof of (1.5).

In Ramanujan's theory of theta-functions, the three theta-functions that play central roles are defined by

$$\varphi(q) := \sum_{n=-\infty}^{\infty} q^{n^2} = \frac{(-q; q)_{\infty}}{(q; -q)_{\infty}}, \quad (1.6)$$

$$\psi(q) := \sum_{n=0}^{\infty} q^{n(n+1)/2} = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}}, \quad (1.7)$$

and

$$f(-q) := \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = (q; q)_{\infty}, \quad (1.8)$$

where

$$(a; q)_{\infty} := \prod_{n=0}^{\infty} (1 - aq^n), \quad |q| < 1.$$

In Chapter 16 of his second notebook [18], Ramanujan records many transformation formulas for $\varphi(q)$, $\psi(q)$ and $f(-q)$. Four of the most important transformation formulas are given by

$$\sqrt{\alpha}\varphi(e^{-\alpha^2}) = \sqrt{\beta}\varphi(e^{-\beta^2}), \quad \alpha\beta = \pi, \quad (1.9)$$

$$2\sqrt{\alpha}\psi(e^{-2\alpha^2}) = \sqrt{\beta}e^{\alpha^2/4}\varphi(-e^{-\beta^2}), \quad \alpha\beta = \pi, \quad (1.10)$$

$$e^{-\alpha/12}\sqrt[4]{\alpha}f(-e^{-2\alpha}) = e^{-\beta/12}\sqrt[4]{\beta}f(e^{-2\beta}), \quad \alpha\beta = \pi^2, \quad (1.11)$$

and

$$e^{-\alpha/24}\sqrt[4]{\alpha}f(e^{-\alpha}) = e^{-\beta/24}\sqrt[4]{\beta}f(e^{-\beta}), \quad \alpha\beta = \pi^2. \quad (1.12)$$

In Section 3, we will prove the transformation formula

$$\sqrt[4]{\alpha}e^{-\alpha/8}\psi(-e^{-\alpha}) = e^{-\beta/8}\sqrt[4]{\beta}\psi(-e^{-\beta}). \quad (1.13)$$

On page 204 of his second notebook [18, p. 204], Ramanujan claims that

$$\left(\frac{\sqrt{5}+1}{2} + R(e^{-2\pi\alpha})\right) \left(\frac{\sqrt{5}+1}{2} + R(e^{-2\pi\beta})\right) = \frac{5+\sqrt{5}}{2} \quad (1.14)$$

and

$$\left(\frac{\sqrt{5}-1}{2} - R(-e^{-2\pi\alpha})\right) \left(\frac{\sqrt{5}-1}{2} - R(-e^{-2\pi\beta})\right) = \frac{5-\sqrt{5}}{2} \quad (1.15)$$

Identities (1.14) and (1.15) were first proved by G. N. Watson [21]. H. H. Chan [12] has proved several identities for $V(q)$ which are similar to (1.14) and (1.15). In Section 4, we will establish three reciprocity theorems for $V(q)$ which are also similar to (1.14) and (1.15).

Ramanujan has recorded many modular equations in his notebooks [5, pp. 204-237], [6, pp. 156-160] which are very useful in the computation of class invariants and the values of theta-functions. In the literature not much attention has been given to find the values of $\psi(q)$ and $\varphi(q)$. But Ramanujan recorded several values of $\varphi(q)$ in his notebooks. For example

$$\varphi(e^{-\pi}) = \frac{\pi^{1/4}}{\Gamma(3/4)},$$

$$\psi(e^{-\pi}) = 2^{-5/8}e^{\pi/8}\frac{\pi^{1/4}}{\Gamma(3/4)},$$

$$\frac{\varphi(e^{-\pi})}{\phi(e^{-3\pi})} = \sqrt[4]{6\sqrt{3} - 9},$$

and

$$\frac{\varphi(e^{-\pi})}{\varphi(e^{-5\pi})} = \sqrt{5\sqrt{5} - 10}.$$

J. M. Borwein and P. B. Borwein [11] first observed that class invariants could be used to calculate certain values of $\varphi(e^{-n\pi})$. Bruce C. Berndt [6] has verified all values for $\varphi(e^{-n\pi})$ claimed by Ramanujan by combining Ramanujan's class invariants with modular equations.

In Section 5, we derive some modular equations and briefly discuss evaluations of theta-functions $\psi(q)$ and $\varphi(q)$. At the end of this section we compute some interesting new explicit evaluations of $V(q)$. Our work is sequel to the works of B. C. Berndt et. al. [7], H. H. Chan [12], K. Ramachandra [16], K. G. Ramanathan [17] and C. Adiga, K. R. Vasuki and M. S. Mahadeva Naika [2]. In this paper we adopt existing methods in the literature and work with $\psi(q)$ instead of $\varphi(q)$ as is done in [12].

2. Integral representations for $V(q)$ and modular equation relating $V(q)$ with $V(q^3)$

Theorem 2.1. *We have*

$$V(q) = \frac{1}{\sqrt[3]{-1 + 9 \exp\left(\int_q^1 \varphi^2(-t) \varphi^2(-t^3) \frac{dt}{t}\right)}} \quad (2.1)$$

$$= \frac{1}{2} \sqrt[3]{1 - \exp\left(-8 \int_0^q \psi^2(t) \psi^2(t^3) dt\right)}. \quad (2.2)$$

Proof of (2.1). Let $F(q) := \frac{\psi^4(q)}{q\psi^4(q^3)}$. Using (1.7) and then taking logarithm on both sides, we find that

$$\begin{aligned} \log F(q) &= 4 \sum_{n=1}^{\infty} \{\log(1 - q^{2n}) - \log(1 - q^{2n-1})\} \\ &\quad + 4 \sum_{n=1}^{\infty} \{\log(1 - q^{3(2n-1)}) - \log(1 - q^{6n})\} - \log q. \end{aligned}$$

Taking derivative on both sides of the above identity, we deduce that

$$\frac{d}{dq} [\log F(q)] = -\frac{1}{q} \left[1 + 4 \sum_{n=1}^{\infty} \frac{(-1)^n n q^n}{1 - q^n} - 12 \sum_{n=1}^{\infty} \frac{(-1)^n n q^{3n}}{1 - q^{3n}} \right].$$

Using Entry 3 (iv) of Chapter 19 of Ramanujan's notebooks [4, p. 226], we find that

$$\frac{d}{dq}[\log F(q)] = -\frac{\varphi^2(-q)\varphi^2(-q^3)}{q}$$

Integrating the above identity over $[q, 1]$ on both sides, we obtain

$$\log F(q) = \int_q^1 \varphi^2(-t)\varphi^2(-t^3)\frac{dt}{t} + \log 9. \quad (2.3)$$

Using Entry 1 (i) of Chapter 20 of Ramanujan's notebooks [4, p. 345] in (2.3) and then exponentiating, we obtain (2.1).

Proof of (2.2). Let $H(q) := \frac{\varphi^4(-q)}{\varphi^4(-q^3)}$. Using (1.6) and then taking logarithm on both sides, we find that

$$\begin{aligned} \log H(q) &= 4 \sum_{n=1}^{\infty} \{\log(1 - q^n) - \log(1 + q^n)\} \\ &\quad + 4 \sum_{n=1}^{\infty} \{\log(1 + q^{3n}) - \log(1 - q^{3n})\}. \end{aligned}$$

Taking derivative on both sides of the above identity, we see that

$$\frac{d}{dq}[\log H(q)] = -\frac{8}{q} \left[\sum_{n=1}^{\infty} \frac{nq^n}{1 - q^{2n}} - 3 \sum_{n=1}^{\infty} \frac{nq^{3n}}{1 - q^{6n}} \right].$$

Using Entry 1 (iii) of Chapter 19 of Ramanujan's notebooks [4, p. 225], we deduce that

$$\frac{d}{dq}[\log H(q)] = -8\psi^2(q)\psi^2(q^3).$$

Integrating the above identity over $[0, q]$ on both sides and then exponentiating, we obtain

$$H(q) = \exp \left(-8 \int_0^q \psi^2(t)\psi^2(t^3)dt \right). \quad (2.4)$$

We deduce (2.2) on employing the following identity [4, p. 345] in (2.4):

$$1 - 8V^3(q) = \frac{\psi^4(-q)}{\psi^4(-q^3)}. \quad (2.5)$$

Theorem 2.2. *Let $V(q)$ be defined as in (1.4). Then*

$$V^3(q) = V^3(q) \frac{1 - V(q^3) + V^2(q^3)}{1 + 2V(q^3) + 4V^2(q^3)}. \quad (2.6)$$

Proof. To prove this theorem we require the following identities [4, p. 345, Entry 1 (i), (ii)]:

$$1 + \frac{1}{V(q^3)} = \frac{\psi(q)}{q\psi(q^9)}, \quad (2.7)$$

$$1 + \frac{1}{V^3(q)} = \frac{\psi^4(q)}{q\psi^4(q^3)}, \quad (2.8)$$

and

$$1 - 3q \frac{\psi(q^9)}{\psi(q)} = \left[1 - 9q \frac{\psi^4(q^3)}{\psi^4(q)} \right]^{1/3}. \quad (2.9)$$

Using (2.7) and (2.8) in (2.9), we obtain

$$1 - \frac{3V(q^3)}{1 + V(q^3)} = \left[1 - \frac{9V^3(q)}{1 + V^3(q)} \right]^{1/3}. \quad (2.10)$$

Cubing both sides of the above identity (2.10), we deduce (2.6).

3. Transformation formula for ψ

In the following theorem we prove a transformation formula for ψ akin to Ramanujan's transformation formulas (1.9)-(1.12).

Theorem 3.1. *If $\alpha\beta = \pi^2$, then*

$$\sqrt[4]{\alpha} e^{-\alpha/8} \psi(-e^{-\alpha}) = e^{-\beta/8} \sqrt[4]{\beta} \psi(-e^{-\beta}). \quad (3.1)$$

Proof. Interchanging α and β in (1.10), we find that

$$2\sqrt{\beta} \psi(e^{-2\beta^2}) = \sqrt{\alpha} e^{\beta^2/4} \varphi(-e^{-\alpha^2}). \quad (3.2)$$

Using (1.10) and (3.2), we deduce that

$$\frac{\psi(e^{-2\alpha^2})}{\psi(e^{-2\beta^2})} = \frac{\beta e^{\alpha^2/4}}{\alpha e^{\beta^2/4}} \frac{\varphi(-e^{-\beta^2})}{\varphi(-e^{-\alpha^2})}, \quad \alpha\beta = \pi. \quad (3.3)$$

Replacing α by $\sqrt{\alpha}$ and β by $\sqrt{\beta}$ in (3.3), we find that

$$\frac{\psi(e^{-2\alpha})\psi(-e^{-\alpha})}{\psi(e^{-2\beta})\psi(-e^{-\beta})} = \sqrt{\frac{\beta}{\alpha}} \frac{e^{\alpha/4}}{e^{\beta/4}}, \quad \alpha\beta = \pi^2. \quad (3.4)$$

Using Entry 25 (iv) of Chapter 16 of Ramanujan's notebooks [4, p. 40] in (3.4), we obtain (3.1).

Remark. The transformation formula (3.1) can also be proved by using the identities (1.11) and (1.12).

4. Reciprocity theorems for $V(q)$

Theorem 4.1. *We have*

(i) *If $\alpha\beta = 1$, then*

$$\left[1 + \frac{1}{V(-e^{-\pi\alpha})}\right] \left[1 + \frac{1}{V(-e^{-\pi\beta})}\right] = 3. \quad (4.1)$$

(ii) *If $3\alpha\beta = 1$, then*

$$\left[1 + \frac{1}{V^3(-e^{-\pi\alpha})}\right] \left[1 + \frac{1}{V^3(-e^{-\pi\beta})}\right] = 9. \quad (4.2)$$

(iii) *If $3\alpha\beta = 1$, then*

$$\left[1 + \frac{1}{V^3(-e^{-\sqrt{2}\pi\alpha})}\right] \left[1 - 8V^3(e^{-\sqrt{2}\pi\beta})\right] = 9. \quad (4.3)$$

Proof of (4.1). Using (2.7), we find that

$$\left[1 + \frac{1}{V(-e^{-\pi\alpha})}\right] \left[1 + \frac{1}{V(-e^{-\pi\beta})}\right] = \frac{\psi(-e^{-\pi\alpha/3})\psi(e^{-\pi\beta/3})}{e^{(-\pi/3)(\alpha+\beta)}\psi(-e^{-3\pi\alpha})\psi(-e^{-3\pi\beta})}. \quad (4.4)$$

From the transformation formula (3.1), we deduce that

$$\frac{\psi(-e^{-\pi\alpha/3})}{e^{(\pi/8)((\alpha/3)-3\beta)}\psi(-e^{-3\pi\beta})} = \sqrt[4]{\frac{9\beta}{\alpha}}. \quad (4.5)$$

Interchanging α and β in (4.5), we obtain

$$\frac{\psi(-e^{-\pi\beta/3})}{e^{(\pi/8)((\beta/3)-3\alpha)}\psi(-e^{-3\pi\alpha})} = \sqrt[4]{\frac{9\alpha}{\beta}}. \quad (4.6)$$

Using (4.5) and (4.6) in (4.4), we obtain (4.1).

Proof of (4.2). Using (2.8), we find that

$$\left[1 + \frac{1}{V^3(-e^{-\pi\alpha})}\right] \left[1 + \frac{1}{V^3(-e^{-\pi\beta})}\right] = \frac{\psi^4(-e^{-\pi\alpha})\psi^4(e^{-\pi\beta})}{e^{-\pi(\alpha+\beta)}\psi^4(-e^{-3\pi\alpha})\psi^4(-e^{-3\pi\beta})}. \quad (4.7)$$

From the transformation formula (3.1), we deduce that

$$\frac{\psi^4(-e^{-\pi\alpha})}{e^{(\pi/2)(\alpha-3\beta)}\psi^4(-e^{-3\pi\beta})} = \frac{3\beta}{\alpha}. \quad (4.8)$$

Interchanging α and β in (4.8), we obtain

$$\frac{\psi^4(-e^{-\pi\beta})}{e^{(\pi/2)(\beta-3\alpha)}\psi^4(-e^{-3\pi\alpha})} = \frac{3\alpha}{\beta}. \quad (4.9)$$

Using (4.8) and (4.9) in (4.7), we obtain (4.2).

Proof of (4.3). Using (2.8), we find that

$$\left[1 + \frac{1}{V^3(-e^{-\sqrt{2}\pi\alpha})}\right] = \frac{\psi^4(e^{-\sqrt{2}\pi\alpha})}{e^{-\sqrt{2}\pi\alpha}\psi^4(e^{-3\sqrt{2}\pi\alpha})} \quad (4.10)$$

Using (1.10) in (4.10), we deduce that

$$\left[1 + \frac{1}{V^3(e^{-\sqrt{2}\pi\alpha})}\right] = 9 \frac{\psi^4(-e^{-\sqrt{2}\pi/\alpha})}{\psi^4(-e^{-\sqrt{2}\pi/3\alpha})} \quad (4.11)$$

Using (2.5) in (4.11), we find that

$$\left[1 + \frac{1}{V^3(-e^{-\sqrt{2}\pi\alpha})}\right] = \frac{9}{1 - 8V^3(e^{-\sqrt{2}\pi/3\alpha})} = \frac{9}{1 - 8V^3(e^{-\sqrt{2}\pi\beta})}.$$

Hence, we complete the proof.

5. Modular equations and evaluations of theta-function and $V(q)$

Theorem 5.1. *Let*

$$P = \frac{\psi(-q)}{q^{1/4}\psi(-q^3)} \quad \text{and} \quad Q = \frac{\varphi(q)}{\varphi(q^3)}.$$

Then

$$Q^4 + P^4Q^4 = 9 + P^4. \quad (5.1)$$

Proof. Using Entry 1 (i) of Chapter 20 of Ramanujan's notebooks [4, p. 345] and (2.5), we find that

$$\frac{\psi^4(q)}{q\psi^4(q^3)} + \frac{\varphi^4(-q)}{\varphi^4(-q^3)} = 9 + \frac{\psi^4(q)\varphi^4(-q)}{q\psi^4(q^3)\varphi^4(-q^3)}.$$

Replacing q by $-q$ in the above identity, we obtain (5.1).

Theorem 5.2. *Let*

$$P = \frac{\psi(-q)}{q\psi(-q^9)} \quad \text{and} \quad Q = \frac{\varphi(q)}{\varphi(q^9)}.$$

Then

$$Q + PQ = 3 + P \quad (5.2)$$

Proof. Using Entry 1 (i) and (ii) of Chapter 20 of Ramanujan's notebooks [4, p. 345], we deduce that

$$\frac{\psi(q^{1/3})}{q^{1/3}\psi(q^3)} + \frac{\varphi(-q^{1/3})}{\varphi(-q^3)} = 3 + \frac{\psi(q^{1/3})\varphi(-q^{1/3})}{q^{1/3}\psi(q^3)\varphi(-q^3)}.$$

Replacing q by $-q^3$ in the above identity, we obtain (5.2).

Theorem 5.3. *Let*

$$P = \frac{\psi(-q)}{q^{1/2}\psi(-q^5)} \quad \text{and} \quad Q = \frac{\varphi(q)}{\varphi(q^5)}.$$

Then

$$Q^2 + P^2Q^2 = 5 + P^2. \quad (5.3)$$

Proof. Changing q to $-q$ in Entry 9 (iii) of Chapter 19 of Ramanujan's notebooks [4, p. 258], we find that

$$\frac{\varphi^2(-q)}{\varphi^2(-q^5)} = 1 - 4W, \quad (5.4)$$

(5.4) where

$$W = \frac{q\chi(-q)f(q^5)f(-q^{20})}{\varphi^2(-q^5)}.$$

Using Entry 9 (vii) of Chapter 19 of Ramanujan's notebooks [4, p. 258], Entry 10 (v) of Chapter 19 of Ramanujan's notebooks [4, p. 262] can be written as

$$\frac{\psi^2(q)}{q\psi^2(q^5)} = 1 + \frac{\varphi(-q^5)f(-q^5)}{q\chi(-q)\psi^2(q^5)}. \quad (5.5)$$

The above identity can be written as

$$\frac{\psi^2(q)}{q\psi^2(q^5)} = 1 + \frac{1}{W}. \quad (5.6)$$

Using (5.4) and (5.6), we find that

$$\frac{\psi^2(q)}{q\psi^2(q^5)} + \frac{\varphi^2(-q)}{\varphi^2(-q^5)} = 5 + \frac{\psi^2(q)\varphi^2(-q)}{q\psi^2(q^5)\varphi^2(-q^5)}.$$

Changing q to $-q$ in the above identity, we obtain (5.3).

Theorem 5.4. *We have*

(i)

$$\frac{\psi(-e^{-\pi/\sqrt{5}})}{e^{-\pi/2\sqrt{5}}\psi(-e^{-\sqrt{5}\pi})} = 5^{1/4}, \quad (5.7)$$

(ii)

$$\frac{\psi(-e^{-3\pi/\sqrt{5}})}{e^{-3\pi/2\sqrt{5}}\psi(-e^{-3\sqrt{5}\pi})} = \frac{5^{1/4}(\sqrt{3}+1)(\sqrt{5}+\sqrt{3})}{2}, \quad (5.8)$$

(iii)

$$\frac{\psi(-e^{-\pi/3\sqrt{5}})}{e^{-\pi/3\sqrt{5}}\psi(-e^{-3\pi/\sqrt{5}})} = \frac{\sqrt{3}(\sqrt{3}-1)(\sqrt{5}-\sqrt{3})}{2}, \quad (5.9)$$

(iv)

$$\frac{\psi(-e^{-\pi/3\sqrt{3}})}{e^{-\pi/12\sqrt{3}}\psi(-e^{-\pi/\sqrt{3}})} = 3^{-1/4}(\sqrt[3]{4} - 1), \quad (5.10)$$

(v)

$$\frac{\psi(-e^{-\pi})}{e^{-\pi/4}\psi(-e^{-3\pi})} = \sqrt[4]{3\sqrt{3}(2 + \sqrt{3})}, \quad (5.11)$$

(vi)

$$\frac{\psi(-e^{-\pi/3\sqrt{5}})}{e^{-\pi/6\sqrt{5}}\psi(-e^{-\sqrt{5}\pi/3})} = \frac{5^{1/4}(\sqrt{3} - 1)(\sqrt{5} - \sqrt{3})}{2}, \quad (5.12)$$

(vii)

$$\frac{\psi(-e^{-\sqrt{5}\pi/3})}{e^{-\sqrt{5}\pi/3}\psi(-e^{-3\sqrt{5}\pi})} = \frac{\sqrt{3}(\sqrt{3} + 1)(\sqrt{5} + \sqrt{3})}{2}, \quad (5.13)$$

(viii)

$$\frac{\psi(-e^{-\pi/\sqrt{3}})}{e^{-\pi/4\sqrt{3}}\psi(-e^{-\sqrt{3}\pi})} = 3^{1/4}, \quad (5.14)$$

(ix)

$$\frac{\psi(-e^{-\pi\sqrt{3}})}{e^{-\pi\sqrt{3}/4}\psi(-e^{-3\sqrt{3}\pi})} = \frac{\sqrt[4]{27}}{\sqrt[3]{4} - 1}, \quad (5.15)$$

(x)

$$\frac{\psi(-e^{-\pi/3})}{e^{-\pi/3}\psi(-e^{-3\pi})} = \sqrt{3}, \quad (5.16)$$

(xi)

$$\frac{\psi(-e^{-\pi/3})}{e^{-\pi/12}\psi(-e^{-\pi})} = \sqrt[4]{\sqrt{3}(2 - \sqrt{3})}, \quad (5.17)$$

(xii)

$$\frac{\psi(-e^{-\pi/\sqrt{3}})}{e^{-\pi/\sqrt{3}}\psi(-e^{-3\sqrt{3}\pi})} = \frac{3}{\sqrt[3]{4} - 1}, \quad (5.18)$$

(xiii)

$$\frac{\psi(-e^{-\pi/3\sqrt{3}})}{e^{-\pi/3\sqrt{3}}\psi(-e^{-\sqrt{3}\pi})} = \sqrt[3]{4} - 1, \quad (5.19)$$

Proofs of the identities (5.7)-(5.19) being similar, for brevity we prove only (5.7)-(5.11).

Proof of (5.7). Putting $\alpha = \frac{\pi}{\sqrt{5}}$ and $\beta = \pi\sqrt{5}$ in the transformation formula (3.1), we obtain (5.7).

Proof of (5.8). By Entry 66 of Chapter 25 of Ramanujan's notebooks [5, p. 233] with q replaced by $-q$, we have

$$P^3Q^3 + 5PQ = Q^4 - 3PQ^3 - 3P^3Q - P^4 \quad (5.20)$$

where

$$P = \frac{\psi(-q)}{q^{1/2}\psi(-q^5)} \quad \text{and} \quad Q = \frac{\psi(-q^3)}{q^{3/2}\psi(-q^{15})}.$$

Using (5.7) in (5.20) with $q = e^{-\pi/\sqrt{5}}$, we find that

$$Q^4 - 5^{1/4}(3 + \sqrt{5})Q^3 - 5^{3/4}(3 + \sqrt{5})Q - 5 = 0. \quad (5.21)$$

Putting $Q = i5^{1/4}T$ in (5.21), we deduce that

$$T^4 + (3 + \sqrt{5})iT^3 - (3 + \sqrt{5})iT - 1 = 0. \quad (5.22)$$

The equation (5.22) can be written as

$$(T^2 - 1)(T^2 + (3 + \sqrt{5})iT + 1) = 0.$$

Solving the above equation, we deduce that

$$T = \pm 1, \quad T = \frac{-(3 + \sqrt{5})i \pm i\sqrt{(3 + \sqrt{5})^2 + 4}}{2}$$

and hence

$$Q = \pm i5^{1/4}, \quad Q = \frac{5^{1/4} \left((3 + \sqrt{5}) \pm \sqrt{(3 + \sqrt{5})^2 + 4} \right)}{2}.$$

Since $Q > 0$, we obtain (5.8).

Proof of (5.9). Putting $q = e^{-\pi/\sqrt{5}}$ in (5.20) and using (5.7), we find that

$$P = 5^{1/4}. \quad (5.23)$$

Also, using (3.1), we see that

$$Q = \frac{\sqrt[4]{45}}{C} \quad (5.24)$$

where

$$C = \frac{\psi(-e^{-\pi/3\sqrt{5}})}{e^{-\pi/3\sqrt{5}}\psi(-e^{-3\pi/\sqrt{5}})}.$$

Using (5.23) and (5.24), we find that

$$PQ = \frac{\sqrt{15}}{C} \quad (5.25)$$

and

$$\frac{P}{Q} = \frac{C}{\sqrt{3}}. \quad (5.26)$$

Using (5.25) and (5.26) in (5.20), we deduce that

$$\sqrt{5} \left[\frac{\sqrt{3}}{C} + \frac{C}{\sqrt{3}} \right] = \left[\frac{\sqrt{3}}{C} + \frac{C}{\sqrt{3}} \right] \left[\frac{\sqrt{3}}{C} - \frac{C}{\sqrt{3}} \right] - 3 \left[\frac{\sqrt{3}}{C} + \frac{C}{\sqrt{3}} \right].$$

Since $\frac{\sqrt{3}}{C} + \frac{C}{\sqrt{3}} \neq 0$, we obtain

$$\frac{\sqrt{3}}{C} - \frac{C}{\sqrt{3}} = \sqrt{5} + 3.$$

Since $C > 0$, solving the above equation we obtain the required result.

Proof of (5.10). Let

$$P = \frac{\psi(-q)}{q^{1/4}\psi(-q^3)} \quad \text{and} \quad Q = \frac{\psi(-q^3)}{q^{3/4}\psi(-q^3)}. \quad (5.27)$$

Using (5.27) in Entry 1 (ii) of Chapter 20 of Ramanujan's notebooks [4, p. 345], we find that

$$Q^3 - P^3Q^2 - 3P^2Q - 3P = 0. \quad (5.28)$$

Putting $q = e^{-\pi/3\sqrt{3}}$ in (5.27), and using the transformation formula (3.1), we see that

$$Q = 3^{1/4}. \quad (5.29)$$

Using (5.29) in (5.28), we deduce that

$$P^3 + 3^{3/4}P^2 + \sqrt{3}P - 3^{1/4} = 0. \quad (5.30)$$

Putting $P = 3^{-1/4}T$ in (5.30), we find that

$$T^3 + 3T^2 + 3T - 3 = 0 \quad (5.31)$$

Putting $x = T + 1$ in (5.31), we obtain

$$x = \sqrt[3]{4}. \quad (5.32)$$

Using (5.32), we obtain the required result.

Proof of (5.11). Putting $q = e^{-\pi/3}$ in (5.28), we find that

$$P = \frac{\psi(-e^{-\pi/3})}{e^{-\pi/12}\psi(-e^{-\pi})} \quad (5.33)$$

and

$$Q = \frac{\psi(-e^{-\pi})}{e^{-\pi/4}\psi(-e^{-3\pi})}. \quad (5.34)$$

Putting $\alpha = \pi/3$ and $\beta = 3\pi$ in (3.1), we deduce that

$$\frac{\psi(-e^{-\pi/3})}{e^{-\pi/4}\psi(-e^{-3\pi})} = \sqrt{3}. \quad (5.35)$$

Using (5.35), we find that

$$PQ = \sqrt{3}. \quad (5.36)$$

Using (5.36) in (5.28), we obtain (5.11).

As the pattern of proof of the following theorem is identical with the proof of Theorem 5.4, we skip the proof.

Theorem 5.5. *We have*

(i)

$$\frac{\varphi(e^{-\pi/\sqrt{5}})}{\varphi(e^{-\sqrt{5}\pi})} = 5^{1/4},$$

(ii)

$$\frac{\varphi(e^{-3\pi/\sqrt{5}})}{\varphi(e^{-3\sqrt{5}\pi})} = \sqrt{\frac{10 + 5\sqrt{3} + 4\sqrt{5} + 2\sqrt{15}}{8 + 5\sqrt{3} + 4\sqrt{5} + 2\sqrt{15}}},$$

(iii)

$$\frac{\varphi(e^{-\pi/3\sqrt{5}})}{\varphi(e^{-3\pi/\sqrt{5}})} = \frac{9 - 3\sqrt{3} + 3\sqrt{5} - \sqrt{15}}{5 - 3\sqrt{3} + 3\sqrt{5} - \sqrt{15}},$$

(iv)

$$\frac{\varphi(e^{-\pi/3\sqrt{3}})}{\varphi(e^{-\pi/\sqrt{3}})} = \sqrt[4]{\frac{27 + (\sqrt[3]{4} - 1)^4}{3 + (\sqrt[3]{4} - 1)^4}},$$

(v)

$$\frac{\varphi(e^{-\pi})}{\varphi(e^{-3\pi})} = \sqrt[4]{6\sqrt{3} - 9},$$

(vi)

$$\frac{\varphi(e^{-\pi/3\sqrt{5}})}{\varphi(e^{-\sqrt{5}\pi/3})} = \sqrt{\frac{10 - 5\sqrt{3} + 4\sqrt{5} - 2\sqrt{15}}{8 - 5\sqrt{3} + 4\sqrt{5} - 2\sqrt{15}}},$$

(vii)

$$\frac{\varphi(e^{-\sqrt{5}\pi/3})}{\varphi(e^{-3\sqrt{5}\pi})} = \frac{9 + 3\sqrt{3} + 3\sqrt{5} + \sqrt{15}}{5 + 3\sqrt{3} + 3\sqrt{5} + \sqrt{15}},$$

(viii)

$$\frac{\varphi(e^{-\pi/\sqrt{3}})}{\varphi(e^{-\sqrt{3}\pi})} = 3^{1/4},$$

(ix)

$$\frac{\varphi(e^{-\pi/\sqrt{3}})}{\varphi(e^{-3\sqrt{3}\pi})} = \sqrt[4]{\frac{27 + 9(\sqrt[3]{4} - 1)^4}{27 + (\sqrt[3]{4} - 1)^4}},$$

(x)

$$\frac{\varphi(e^{-\pi/3})}{\varphi(e^{-3\pi})} = \sqrt{3},$$

(xi)

$$\frac{\varphi(e^{-\pi/3})}{\varphi(e^{-\pi})} = \sqrt[4]{3 + 2\sqrt{3}},$$

(xii)

$$\frac{\varphi(e^{-\pi/\sqrt{3}})}{\varphi(e^{-3\sqrt{3}\pi})} = \frac{3\sqrt[3]{4}}{2 + \sqrt[3]{4}},$$

(xiii)

$$\frac{\varphi(e^{-\pi/3\sqrt{3}})}{\varphi(e^{-\sqrt{3}\pi})} = 4^{-1/3}(2 + \sqrt[3]{4}).$$

Using (2.7) and (2.8) in Theorem 5.4, we obtain the following values of the cubic continued fraction of Ramanujan.

Theorem 5.6. *We have*

$$V(-e^{-\pi\sqrt{5}}) = \frac{(3 - \sqrt{5})(\sqrt{3} - \sqrt{5})}{4},$$

$$V(-e^{-\pi/\sqrt{5}}) = \frac{(\sqrt{3} + \sqrt{5})(\sqrt{5} - 3)}{4},$$

$$V(-e^{-\pi/\sqrt{3}}) = \frac{-1}{\sqrt[3]{4}},$$

$$V(-e^{-\pi\sqrt{3}}) = \frac{1 - \sqrt[3]{4}}{2 + \sqrt[3]{4}},$$

$$V(-e^{-\pi/3\sqrt{3}}) = \frac{-1}{\sqrt[3]{3^{-1}(\sqrt[3]{4} - 1)^4 + 1}},$$

$$V(-e^{-\pi}) = \frac{1 - \sqrt{3}}{2},$$

$$V(-e^{-\pi/3}) = \frac{-1}{\sqrt[3]{2(\sqrt{3} - 1)}}.$$

Remark. A different proof of Theorem 5.6 (i) can be found in [12] and Theorem 5.6 (iii) was first proved by C. Adiga et al. [2]. Other values of V in Theorem 5.6 appear to be new to literature.

Open Problem. Using (2.10) one can express $V(q^3)$ in terms of $V(q)$ which gives a triplication formula for $V(q)$. Triplication formula is important in the development of elliptic functions to alternative bases. The only triplication formula known so far is that of the Borweins. Can one develop a cubic theory associated with $V(q)$ similar to that of Borweins?

Acknowledgement

Authors are thankful to the referee for his useful suggestions and comments. This work was supported by Korea Research Foundation Grant (KRF-2002-050-C00001).

REFERENCES

- [1] C. Adiga, K. R. Vasuki and M. S. Mahadeva Naika, *On some new identities involving integrals of theta-functions*, Advanced Studies in Contemporary Mathematics **3** (2001), 1–11.
- [2] C. Adiga, K. R. Vasuki and M. S. Mahadeva Naika, *Some new explicit evaluations of Ramanujan's cubic continued fraction*, New Zealand Journal of Mathematics (to appear).
- [3] G. E. Andrews, Ramanujan's, "lost" notebook, III. *The Rogers-Ramanujan continued fraction*, Adv. Math. **41** (1981), 186–208.
- [4] B. C. Berndt, *Ramanujan's notebooks, part III*, Springer-Verlag, New York, 1991.
- [5] B. C. Berndt, *Ramanujan's notebooks, part IV*, Springer-Verlag, New York, 1994.
- [6] B. C. Berndt, *Ramanujan's notebooks, part V*, Springer-Verlag, New York, 1998.
- [7] B. C. Berndt, H. H. Chan and L.-C. Zhang, *Ramanujan's class invariants and cubic continued fraction*, Acta Arith. **73** (1995), 67–85.
- [8] S. Bhargava, *On Rogers-Ramanujan identities*, in *Srinivasa Ramanujan (1887-1920)*, Macmillan India Ltd., Madras, 1988, 75–81.
- [9] S. Bhargava and C. Adiga, *On some continued fraction identities of Srinivasa Ramanujan*, Proc. Amer. Math. Soc. **92** (1984), 13–18.
- [10] S. Bhargava and C. Adiga, *Two generalizations of Ramanujan's continued fraction identities*, in *Number Theory*, K. Alladi, ed., Lecture Notes in Math. No. 1122, Springer-Verlag, Berlin, 1985, 56–62.
- [11] J. M. Borwein and P. B. Borwein, *Pi and the AGM*, Wiley, New York, 1987.
- [12] H. H. Chan, *On Ramanujan's cubic continued fraction*, Acta Arith. **73** (1995), 343–355.
- [13] R.Y. Denis, *On certain q -series and continued fractions*, Math. Student **44** (1983), 70–76.
- [14] R.Y. Denis, *On basic hypergeometric functions and continued fractions*, Math. Student **52** (1984), 129–136.
- [15] R.Y. Denis, *On certain summation of q -series and identities of Rogers-Ramanujan type*, J. Math. Phys. Sci. **22** (1988), 87–99.
- [16] K. Ramachandra, *Some applications of Kronecker's limit formulas*, Annals of Mathematics **80** (1964), 104–148.
- [17] K. G. Ramanathan, *On Ramanujan's continued fraction*, Acta Arith. **43** (1984), 209–226.

- [18] S. Ramanujan, *Notebooks (2 volumes)*, Tata Institute of Fundamental Research, Bombay, 1957.
- [19] S. Ramanujan, *The lost notebook and other unpublished papers*, Narosa, New Delhi, 1988.
- [20] Seung Hwan Son, *Some integrals of theta-functions in Ramanujan's 'lost' notebook*, CRM Proceedings and Lecture Notes **19** (1999), 323–332.
- [21] G. N. Watson, *Theorems stated by Ramanujan (ix): two continued fractions*, J. London Math. Soc. **4** (1929), 231–237.